



Hexavalent chromium removal from aqueous solutions using biogenic iron nanoparticles: Kinetics and equilibrium study

P. Prema^a, Van-Huy Nguyen^b, Karthikeyan Venkatachalam^c, J.M. Murugan^d, Hayssam M. Ali^e,
Mohamed Z.M. Salem^f, Balasubramani Ravindran^g, P. Balaji^d  

- ^a Department of Zoology, VHN Senthikumara Nadar College (Autonomous), Virudhunagar, Tamilnadu, India
- ^b Faculty of Biotechnology, Binh Duong University, Thu Dau Mot, Viet Nam
- ^c Faculty of Innovative Agriculture and Fishery Establishment Project, Prince of Songkla University Surat Thani Campus, Makhm Tia, Mueang, Surat Thani, 84000, Thailand
- ^d PG and Research Centre in Biotechnology, MGR College, Hosur, India
- ^e Department of Botany and Microbiology College of Science, King Saud University, P.O. Box 2455, Riyadh, 11451, Saudi Arabia
- ^f Department of Forestry and Wood Technology, Faculty of Agriculture (EL-Shatby), Alexandria University, Alexandria, 21545, Egypt
- ^g Department of Environmental Energy and Engineering, Kyonggi University Youngtong-Gu, Suwon, Gyeonggi-Do, 16227, Republic of Korea